

方知库
Eco-Environmental
Knowledge Web

环境科学

ENVIRONMENTAL SCIENCE

ISSN 0250-3301 CODEN HCKHDV

HUANJING KEXUE

■ 主办 中国科学院生态环境研究中心

■ 出版 科学出版社



2020

Vol.41 No.1

第41卷 第1期

目次

近20年来中国PM _{2.5} 污染演变的时空过程	时燕, 刘瑞梅, 罗毅, 杨昆 (1)
2007~2016年上海颗粒物浓度特征与气候背景异同分析	郑庆峰, 史军, 谈建国, 段玉森, 林燕芬, 徐卫忠 (14)
夏季大气PM _{2.5} 中元素特征及源解析:以华中地区平顶山-随州-武汉为例	刘威杰, 石明明, 程敏, 黎莹, 朱戈昊, 邢新丽, 孔少飞, 张家泉, 祁士华 (23)
呼包鄂地区冬季、夏季PM _{2.5} 污染特征	李伊明, 刘佳媛, 史国良, 皇甫延琦, 张昕, 杨奕, 冯银厂 (31)
2019年元旦前后石家庄市重污染过程PM _{2.5} 污染特征及来源解析	周静博, 段青春, 王建国, 杨鹏, 刘煥武, 李梅, 靳伟 (39)
基于模型模拟的成都市PM _{2.5} 污染来源解析	许艳玲, 易爱华, 薛文博 (50)
北京市PM _{2.5} 中砷污染特征	石绍萱, 杨艳蓉, 秦娟娟, 周雪明, 段青春, 谭吉华, 陈荣志 (57)
夏秋季宁波市气态元素汞(GEM)污染特征及潜在源区贡献分析	易辉, 林佳梅, 蔡秋亮, 王克强, 陈进生, 佟磊, 李建荣, 肖航 (65)
安阳市大气PM _{2.5} 中水溶性离子季节特征及来源解析	孙有昌, 姜楠, 王申博, 段时光, 张瑞芹 (75)
新乡市夏冬季节PM _{2.5} 稳定碳同位素特征分析	闫广轩, 宋鑫, 张靖雯, 张朴真, 王力鹏, 张佳羽, 杨争, 曹治国, 樊静, 王跃思, 刘子锐 (82)
基于颗粒物化学组分粒径分布特征的源解析方法构建与评估	梁永莉, 田瑛泽, 刘童, 冯银厂 (90)
超低排放改造后燃煤电厂细颗粒物排放特征	王润芳, 马大卫, 姜少毅, 朱仁斌, 何军, 张本耀 (98)
京津冀地区城市臭氧污染趋势及原因探讨	余益军, 孟晓艳, 王振, 周崑, 于红霞 (106)
安阳市近地面臭氧污染特征及气象影响因素分析	杨健, 尹沙沙, 于世杰, 王丽娟, 张瑞芹, 郝建尤, 苏方成, 齐艳杰 (115)
长江三角洲2017年机动车IVOCs排放清单构建及其对SOA的生成影响	王倩, 黄凌, 王杨君, 殷司佳, 张琪, 易鑫, 李莉 (125)
华东典型城市非道路移动机械排气烟度	鲁君, 胡馨遥, 杨强, 唐伟, 夏阳, 井宝莉, 陈昀, 黄成 (133)
国内外储油库VOCs排放现状与标准分析	胡玮, 任碧琪, 黄玉虎, 秦建平, 卞吉玮, 吴清柱 (139)
四乙基愈创木酚液相·OH氧化SOA产率及特征分析:初始浓度的影响	庄雨, 陈彦彤, 李旭东, 马帅帅, 赵竹子, 叶招莲, 盖鑫磊 (146)
乌鞘岭南、北坡降水稳定同位素特征及水汽来源对比	徐秀婷, 贾文雄, 朱国锋, 马兴刚, 张昱, 袁瑞丰, 张志远, 石阳 (155)
长江源多年冻土区地下水氢氧稳定同位素特征及其影响因素	汪少勇, 何晓波, 丁永建, 常福宣, 吴锦奎, 胡召富, 王利辉, 杨贵森, 邓明珊 (166)
巴尔喀什湖流域水化学和同位素空间分布及环境特征	沈贝贝, 吴敬禄, 吉力力·阿不都外力, A. S. Saparov, Gulnura Isanova (173)
城市湖泊水环境整治对改善水质的影响:以蠡湖近30年水质变化为例	田伟, 杨周生, 邵克强, 潘辉, 胡洋, 白承荣, 姜星宇, 高光, 汤祥明 (183)
典型药物及个人护理品在黄东海海域水体中的检测、分布规律及其风险评估	陈贤, 张彩杰, 杨桂朋, 张婧, 刘梦 (194)
长江南京段水源水中有机磷酸酯的污染特征与风险评估	李栋, 张圣虎, 张芹, 王博, 卜元卿, 赵欣, 李辉, 宋宁慧, 郭瑞昕 (205)
雄安新区-白洋淀冬季冰封期水体溶解性有机物的空间分布、光谱特征及来源解析	周石磊, 孙悦, 张艺冉, 蒋永丰, 尹璐, 黄廷林, 李再兴, 罗晓, 战江, 王函 (213)
白洋淀表层沉积物元素的空间特征、风险评价及来源分析	汪敬忠, 刘卓, 魏浩, 吴玉会, 占水娥, 朱迟, 张益森 (224)
松花江表层沉积物PAEs分布特征及生态风险评价	王欢, 杨永哲, 王海燕, 董伟丰, 闫国凯, 常洋, 李泽文, 赵远哲, 凌宇 (232)
鄱阳湖湖口-长江段沉积物中微塑料与重金属污染物的赋存关系	李文华, 简敏菲, 刘淑丽, 江玉梅, 邓艳彬, 朱琳 (242)
臭氧对生物活性炭中微生物及出水消毒副产物的影响	刘宝明, 王夕希, 张晓夏, 顾艳芳, 李雨平, 阮文权, 缪恒峰, 潘杨 (253)
兰州市西固区儿童饮用水重金属暴露与健康风险精细化评估	陈月芳, 孙善伟, 段小丽, 许锦荣, 王娟, 康艺瑾, 梁为纲, 曹素珍 (262)
镁铁层状双金属氢氧化物对磷酸盐的吸附作用及对内源磷释放的控制效果及机制	吴俊麟, 林建伟, 詹艳慧, 程永前, 柏晓云, 辛慧敏, 常明玥 (273)
一硫代磷酸盐在介质上的吸附特征及机制	廖丹雪, 单慧媚, 彭三曦, 潘敖然, 黄健, 陈辉, 赵超然 (284)
酸碱复合改性海泡石亚结构特征及其对Cd(II)吸附性能	谢厦, 徐应明, 闫翠侠, 罗文文, 孙约兵 (293)
脱氮菌剂在低溶解性黑臭水体中氮代谢特征	赵志瑞, 刘硕, 李铎, 李海萍, 李方红, 张佳瑶, 白志辉 (304)
宏基因组方法分析医药化工废水厂中抗生素耐药菌及耐药基因	陈红玲, 张兴桃, 王晴, 姚沛琳 (313)
淀粉废水处理系统中活性污泥的微生物群落结构及多样性分析	赵婷婷, 乔凯, 王蕾, 张伟, 竺建荣, 唐一, 胡滨, 王香莲, 杨静, 景在涛 (321)
基于SBR-ABR实现PN-SAD耦合工艺的运行与优化调控	陈重军, 张敏, 姜滢, 郭萌蕾, 谢嘉玮, 谢军祥, 沈耀良 (330)
基于间歇饥饿的SNAD工艺运行	李冬, 刘志诚, 徐贵达, 王玉颖, 张杰 (337)
反硝化-高效部分亚硝化-厌氧氨氧化工艺处理老龄垃圾渗滤液	陈小珍, 汪晓军, Karasuta Chayangkun, 周松伟, 钟中, 陈振国, 陈晓坤 (345)
硝化微颗粒污泥快速培养及其亚硝化功能快速实现	刘文如, 宋家俊, 王建芳, 沈耀良 (353)
高频曝停下曝时间对亚硝化颗粒污泥性能的影响	张杰, 劳会妹, 李冬, 魏子清 (360)
中国东北沈抚地区土壤中PBDEs的残留、来源及风险评价	向鑫鑫, 鲁垠涛, 阮起扬, 赖偲, 孙绍斌, 姚宏, 张战胜 (368)
某焦化场地非均质包气带中多环芳烃(PAHs)来源及垂向分布特征	孟祥帅, 吴萌萌, 陈鸿汉, 岳希, 陶抒远 (377)
北京蔬菜地土壤中抗生素抗性基因与可移动元件的分布特征	张汝凤, 宋渊, 高浩泽, 程首涛, 孙艳梅, 王旭明 (385)
长期施肥和耕作下紫色土坡耕地土壤C、N、P和K化学计量特征	杜映妮, 李天阳, 何丙辉, 贺小容, 付适 (394)
亚热带稻田土壤碳氮磷生态化学计量学特征	宋佳龄, 盛浩, 周萍, 段良霞, 周清, 张杨珠 (403)
宛山荡农田土壤氮迁移过程反硝化与厌氧氨氧化	陈志浩, 覃云斌, 丁帮琛, 胡优优, 陈湜, 李正魁 (412)
秸秆炭化还田对滴灌棉田土壤微生物代谢功能及细菌群落组成的影响	王晶, 马丽娟, 刘泽华, 陆伟, 侯振安 (420)
基于PMF模型的九龙江流域农田土壤重金属来源解析	黄华斌, 林承奇, 胡恭任, 于瑞莲, 郝春莉, 陈枫桦 (430)
塔里木盆地东南缘绿洲区土壤钾空间分布及农作物钾富集特征	陈云飞, 周金龙, 曾妍妍, 王松涛, 杜江岩, 孙英, 顾思博 (438)
广西典型碳酸盐岩区农田土壤-作物系统重金属生物有效性及迁移富集特征	马宏宏, 彭敏, 刘飞, 郭飞, 唐世琪, 刘秀金, 周亚龙, 杨柯, 李括, 杨峥, 成杭新 (449)
不同钝化剂对微碱性土壤镉、镍形态及小麦吸收的影响	张静静, 朱爽阁, 朱利楠, 柳海涛, 杨金康, 化党领 (460)
不同钝化剂对鸡粪堆肥重金属钝化效果及其腐熟度指标的影响	栾润宇, 高珊, 徐应明, 纪艺凝, 闫翠侠, 孙约兵 (469)
黄土丘陵区退耕草地土壤呼吸及其组分对氮磷添加的响应	周俊杰, 陈志飞, 杨全, 赖帅彬, 王智, 徐炳成 (479)
炉渣与生物炭施加对福州平原水稻田温室气体排放的后续影响	蓝兴福, 王晓彤, 周雅心, 金强, 许旭萍, 王维奇 (489)
《环境科学》征订启事(89)	
《环境科学》征稿简则(97)	
信息(303, 329, 344)	

长江南京段水源水中有机磷酸酯的污染特征与风险评估

李栋^{1,2}, 张圣虎², 张芹², 王博², 卜元卿², 赵欣², 李辉³, 宋宁慧^{2*}, 郭瑞昕^{1*}

(1. 中国药科大学工学院, 南京 211198; 2. 生态环境部南京环境科学研究所, 南京 210042; 3. 北京百灵天地环保科技股份有限公司, 北京 100045)

摘要: 为探究长江南京段水源水中有机磷酸酯(organophosphate esters, OPEs)的污染特征、时空分布、生态风险和健康风险, 利用固相萃取-高效液相色谱-串联质谱法测定了13种OPEs。结果表明, 除磷酸三(2,3-二溴丙基)酯外, 其余12种OPEs均有不同程度的检出, 总浓度范围为85.21~1557.96 ng·L⁻¹, 氯代烷基磷酸酯是主要化合物, 其中检出浓度最高的是磷酸三(2-氯乙基)酯[tri(2-chloroethyl) phosphate, TCEP], 高达447.08 ng·L⁻¹。长江南京段水源水中OPEs呈现明显的季节变化特征, 夏季总检出浓度为220~1557.96 ng·L⁻¹, 平均浓度是493.78 ng·L⁻¹, 是春秋两季的1.7~2.6倍。生态风险评估显示磷酸三甲苯酯和2-乙基己基二苯基磷酸酯对有机体(藻类, 甲壳类动物和鱼类)具有中或高等风险。高暴露浓度下, OPEs的总非致癌风险为4.41×10⁻³~2.91×10⁻², 均小于1, 0~3个月的婴儿最高, 总致癌风险值为5.88×10⁻⁷~3.89×10⁻⁶, 其中TCEP和磷酸三(1,3-二氯-2-丙基)酯对儿童有潜在的致癌风险。长江南京段水源水中OPEs对儿童的长期暴露风险需引起高度重视。

关键词: 长江南京段; 水源水; 有机磷酸酯(OPEs); 污染特征; 季节特征; 生态风险; 健康风险

中图分类号: X522; X820.4 文献标识码: A 文章编号: 0250-3301(2020)01-0205-08 DOI: 10.13227/j.hjxx.201907149

Occurrence and Risk Assessment of Organophosphate Esters in Source Water of the Nanjing Section of the Yangtze River

LI Dong^{1,2}, ZHANG Sheng-hu², ZHANG Qian², WANG Bo², PU Yuan-qing², ZHAO Xin², LI Hui³, SONG Ning-hui^{2*}, GUO Rui-xin^{1*}

(1. School of Engineering, China Pharmaceutical University, Nanjing 211198, China; 2. Nanjing Institute of Environmental Sciences, Ministry of Ecology and Environment, Nanjing 210042, China; 3. Beijing Larkworld Environmental Technology Incorporated Company, Beijing 100045, China)

Abstract: We explored the pollution characteristics, spatial and temporal distribution, ecological risk, and human health risk of organophosphate esters (OPEs) in the source water of the Nanjing section of the Yangtze River. Thirteen OPEs were determined by solid phase extraction-high performance liquid chromatography-tandem mass spectrometry. The results showed that twelve OPEs were detected to different extents with total concentrations ranging from 85.21 to 1557.96 ng·L⁻¹ [excluding tris (2,3-dibromopropyl) phosphate]. Chloroalkyl phosphate was the main compound and tris(2-chloroethyl) phosphate (TCEP) was the most abundant of the thirteen OPEs, which reached up to 447.08 ng·L⁻¹. Seasonal variation of OPEs showed that the concentrations of OPEs in summer ranged from 220-1557.96 ng·L⁻¹, with the average concentration of 493.78 ng·L⁻¹ being 1.7-2.6 times higher than of that in spring and autumn. An ecological risk assessment showed that tricresyl phosphate and 2-ethylhexyl diphenyl phosphate were associated with moderate or high risks to organisms (algae, crustaceans, and fish). At a high exposure concentration, the total non-carcinogenic risk of OPEs in source water ranged from 4.41×10⁻³-2.91×10⁻², with the highest risk being associated with children aged 0-3 months. The total cancer risk value was 5.88×10⁻⁷-3.89×10⁻⁶, among which TCEP and Tris (1,3-dichloro-2-propyl) phosphate was associated with a potential risk for children. We conclude that the long-term exposure risk for children of OPEs in the source water of the Nanjing section of the Yangtze River deserve more attention.

Key words: Nanjing section of the Yangtze River; source water; organophosphate esters (OPEs); pollution characteristics; seasonal variation; ecological risk; health risk

有机磷酸酯(Organophosphate esters, OPEs), 因其良好的阻燃、润滑和增塑效果, 成为多溴联苯醚等溴代阻燃剂的主要替代品, 广泛应用于塑料制品、建筑材料、家装饰品、电子产品和纺织品等中^[1~3], 其用量近几年大幅增长, 1992年全球OPEs消费量为10.2万t, 2013年增加到37万t, 2015年数据显示已经增至68万t^[4~6]。

由于OPEs主要以非化学键方式添加进产品, 在生产、使用和废弃过程中很容易通过挥发、磨损和渗出等方式进入水体^[7~9], 而现阶段水处理技术

对OPEs的去除效果并不明显^[3,10,11], 因此很容易引起水环境污染。目前, 我国渤海^[12]、太湖^[13]、骆马湖^[14]、松花江^[15]和珠江^[16]等水体中均检出了OPEs, 而毒理研究表明OPEs具有内分泌干扰、致癌和致基因突变等多种毒性^[3,5], 潜在影响饮用水

收稿日期: 2019-07-17; 修订日期: 2019-08-19

基金项目: 国家自然科学基金项目(21507036); 国家重点研发计划项目(2017YFD0800705); 云南省环境保护厅项目(YNZDZB[2017]002)

作者简介: 李栋(1996~), 女, 硕士研究生, 主要研究方向为污染物环境归趋与风险评估, E-mail: dongli0619@163.com

* 通信作者, E-mail: Snh@nies.org; grxcpu@163.com

源地水质安全.

长江是南京居民主要饮用水源,水质安全格外重要,但关于其有机磷酸酯的污染特征和风险评估鲜有报道.本研究通过分析长江南京段水源水中 OPEs 的浓度以及不同时期(丰水期、枯水期和平水期)水体中污染状况,揭示 OPEs 的季节变化特征,并评估在不同人群中 OPEs 对生态和人体健康的潜在风险,以期为城乡饮用水安全和 OPEs 的风险控制提供参考依据.

1 材料与方法

1.1 仪器与试剂

高效液相色谱-串联质谱仪(HPLC-Agilent Technologies 1290 Infinity,美国 MS-ABSCIEX QTRAP 4500);旋转蒸发仪(R-300,瑞士 BUCHI 公司);萃取小柱[Oasis HLB, 6 mL·(500 mg)⁻¹,美国 Waters 公司];高速离心机(Centrifuge 5430R,德国 Eppendorf 公司);超纯水仪(Milli-Q,美国 Millipore 公司);电子天平(AG-285,瑞士 Mettler 公司).

13 种 OPEs 标准品,包括 6 种烷基磷酸酯:磷酸三(2-丁氧基乙基)酯[tri(2-butoxyethyl) phosphate, TBOEP, 95.0%];磷酸三(2-乙基己基)酯[tri(2-ethylhexyl) phosphate, TEHP, 98.0%];磷酸三乙酯(triethyl phosphate, TEP, 99.0%);磷酸三甲酯(trimethyl phosphate, TMP, 99.0%);磷酸三正丁酯(tri-n-butyl phosphate, TnBP, 99.0%)和磷酸三丙酯(tripropyl phosphate, TPrP, 99.0%);4 种卤代烷基磷酸酯:磷酸三(2-氯乙基)酯[tri(2-chloroethyl) phosphate, TCEP, 97.0%];磷酸三(2-氯异丙基)酯[tri(2-chloropropyl) phosphate, TCPP, 98.0%];磷酸三(2,3-二溴丙基)酯[tri(2,3-dibromopropyl) phosphate, TDBPP, 98.0%]和磷酸三(1,3-二氯-2-丙基)酯[tri(1,3-dichloropropyl) phosphate, TDCPP, 95.0%];和 3 种芳香族磷酸酯:2-乙基己基二苯基磷酸酯(2-ethylhexyl diphenyl phosphate, EHDPP, 95.0%);磷酸三甲苯酯(tricresyl phosphate, TCrP, 95.0%)和磷酸三苯酯(triphenyl phosphate, TPhP, 99.0%)均购于百灵威科技有限公司.替代物 TnBP-d₂₇ 和内标物 TPhP-d₁₅ 购于 Cambridge Isotope Laboratories(Tewksbury, MA, USA).甲醇和乙酸乙酯(色谱纯)购于德国 Merck 公司,甲酸(色谱纯)购自国药集团药业股份有限公司.

1.2 样品采集

分别于 2018 年 8 月(夏季,丰水期)、2018 年 11 月(秋季,枯水期)和 2019 年 3 月(春季,平水期),采集长江南京段水源水表层水样,选取 16 个

采样点(S1~S16),位置见图 1.使用有机玻璃水样采集器采集水样,采集深度为 0~1 m,每个样点采集 2 L,置于棕色磨口塞玻璃瓶中,尽快运回实验室,4℃避光保存,24 h 内进行前处理.




图 1 长江南京段水源水采样点示意

Fig. 1 Sampling sites of source water in the Nanjing section of the Yangtze River

1.3 样品前处理

参照并优化本课题组建立的方法^[17],准确移取 500 mL 经 0.45 μm 玻璃纤维膜过滤后的水样,加入 10 ng 的替代物 TnBP-d₂₇ 以校正预处理过程中目标物的损失.之后用固相萃取小柱进行富集.预先依次用 5 mL 乙酸乙酯、5 mL 甲醇和 5 mL 超纯水对 HLB 小柱进行活化,上样后,用 5 mL 超纯水淋洗 HLB 小柱,然后在真空负压条件下抽干 30 min 至小柱充分干燥;再用 5 mL 乙酸乙酯以 1 mL·L⁻¹ 的流速洗脱 2 次.收集的洗脱液于 40℃ 下经旋转蒸发浓缩至干,最后用甲醇复溶至 1 mL,在 HPLC-MS/MS 分析前加入 10 ng 的内标物 TPhP-d₁₅ 以便后续进行定量分析.

1.4 仪器分析

高效液相测定条件:采用 ZORBAX Eclipse Plus C₁₈ 色谱柱(150 mm×2.1 mm, 3.5 μm,美国 Agilent 公司);柱温设置为 30℃;以体积分数为 0.2% 的甲酸(A)和乙腈(B)作为流动相,梯度洗脱.程序为:0~8 min, 30% A;8~8.1 min, 30%~5% A;8.1~16 min, 5.0% A;16~20 min, 5%~30% A;流速为 0.3 mL·min⁻¹,进样体积为 5 μL.

质谱检测条件:采用电喷雾离子源(ESI)、正

离子 MRM 监测模式;离子源温度为 500℃,喷雾电 压力分别为 35.0、55.0 和 60.0 kPa,其他参数见
压设置为5 500 V;气帘气,喷雾气和辅助加热气的 表 1.

表 1 目标物的主要参数¹⁾
Table 1 Parameters of target compounds

化合物	母离子(<i>m/z</i>)	子离子(<i>m/z</i>)	碰撞能/eV	去簇电压/V	入口电压/V	出口电压/V
TMP	140.9	108.8 *	23	62	10	7
		79.1	28	62	10	7
TEP	182.6	98.7 *	23	50	10	7
		126.7	15	50	10	7
TPrP	225.1	98.9 *	23	93	10	10
		141.1	13	102	10	10
TnBP	267.1	98.9 *	10	110	10	7
		155	20	110	10	7
TBOEP	399.2	198.9 *	21	73	10	7
		298.9	17	73	10	7
TCPP	327	99.0 *	32	63	7	10
		174.7	16	63	7	10
TDCPP	430.9	99.0 *	33	82	7	10
		208.7	21	86	7	10
TPhP	327.1	214.9 *	34	114	10	10
		152.1	55	114	10	10
TCrP	369.3	164.8 *	55	133	10	10
		243.3	55	133	10	10
TCEP	284.8	223.1 *	15	79	10	10
		99.3	31	79	10	10
TEHP	435.5	98.7 *	43	70	5	10
		112.9	12	70	5	10
TDBPP	698.5	299.1 *	21	102	10	10
		500.8	21	102	10	10
EHDPP	362.7	250.9 *	12	78	10	7
		151.9	45	78	7	10
TnBP-d ₂₇	294.4	102.0 *	17	112	10	15
		166.1	25	125	10	15
		230.1	44	110	10	7
TPhP-d ₁₅	342.2	160.0 *	48	110	10	10
		223.1	36	110	10	10
		262	36	110	10	10

1) * 表示定量离子

1.5 质量控制与保证(QA/QC)

为保证结果的准确性,在取样、运输、储存和提取过程中避免使用塑料和橡胶材料,并依次进样溶剂、标准品和程序空白以检查背景污染.使用内标法定量,标准曲线在 0.1 ~ 500 mg·L⁻¹ 范围内线性良好,相关系数(*R*²)均大于 95%.以信噪比 *S*/*N* = 10 计算定量限(LOQ),13 种 OPEs 的 LOQ 为 0.37 ~ 265.33 ng·L⁻¹.在 100 mL 超纯水添加 5 μg·L⁻¹ 的 13 种 OPEs 混合标准溶液,同时设置空白对照试验,各设置 3 个平行样,各目标化合物的回收率为 66.4% ~ 106.9%,相对标准偏差为 0.38% ~

14.16%.

1.6 风险评价

1.6.1 生态风险评估

OPEs 的生态风险常用风险熵(RQ)评估,计算公式为^[8,18]:

$$RQ = MEC/PNEC \tag{1}$$

式中,MEC 为环境实际测量浓度,PNEC 为无效应浓度,通常由急性和慢性毒性数据(半数致死浓度 LC₅₀、半数效应浓度 EC₅₀、最大无效应浓度 NOEC 等)除以评估因子(AF)得到^[8,13];当 RQ < 0.1 时,风险较小或可忽略不计,当 0.1 ≤ RQ < 1.0 时具有

中等风险,而当 $RQ \geq 1.0$ 时,则认为具有高风险。

1.6.2 人体健康风险评估

采用美国环保署(US EPA)推荐的健康风险模型^[19,20],评价长江南京段水源水中 OPEs 日摄入量,并通过致癌风险和非致癌风险来分析 OPEs 对人体健康的影响。

OPEs 的日均暴露量(ADD)按下式计算^[10,11]:

$$ADD = c \times IR \times AP/BW \quad (2)$$

式中, c 为 OPEs 的浓度($\text{ng} \cdot \text{L}^{-1}$); IR 为日均饮水量($\text{L} \cdot \text{d}^{-1}$); AP 为饮水的吸收率, AP 取 100%; BW 为平均体重(kg)。中国人群暴露参数手册(成人卷)^[21]提供了江苏成人平均体重 BW 为 63.2 kg,全年日均饮水量 IR 为 $2.33 \text{ L} \cdot \text{d}^{-1}$,其他参数采用 US EPA 和相关文献推荐值^[22~24]。

非致癌风险用危害商(HQ)表征,计算公式如下^[10,13]:

$$HQ = ADD/RfD \quad (3)$$

式中, RfD 为 USPA 建议的参考剂量;当 $HQ < 1$ 时,非致癌风险较小或忽略不计;当 $HQ > 1$ 时,则表明具有非致癌风险。对多污染物暴露情形,非致癌总风险即为各 OPEs 危害商 HQ 之和。

致癌风险(CR)是致癌物质的暴露剂量乘以相应的致癌斜率因子(SFO)得到,计算如下^[11,14,20]:

$$CR = ADD \times SFO \quad (4)$$

当 CR 低于 10^{-6} ,致癌风险可以忽略不计,当 CR 介于 $10^{-6} \sim 10^{-4}$ 之间,表明具有潜在的致癌风险,若 CR 值高于 10^{-4} ,则认为具有高风险,各种致癌目标物致癌风险之和即为“总致癌风险”。相关参数见表 2。

表 2 健康风险计算参数

年龄段	IR / $\text{L} \cdot \text{d}^{-1}$	BW / kg	目标物	RfD / $\text{ng} \cdot (\text{kg} \cdot \text{d})^{-1}$	SFO / $\text{kg} \cdot \text{d} \cdot \text{ng}^{-1}$
0~3 个月	1.15	5.60	TMP	0.010	0.020
3~6 个月	1.14	7.20	TEP	0.125	—
6~12 个月	1.18	9.40	TnBP	0.010	0.009
1~2 岁	0.85	12.0	TBOEP	0.015	—
2~3 岁	0.83	13.8	TCPP	0.010	—
3~6 岁	1.16	19.0	TDCPP	0.020	0.031
6~11 岁	1.55	36.0	TPhP	0.070	—
11~16 岁	1.9	56.0	TCrP	0.013	—
16~18 岁	1.77	57.0	TCEP	0.007	0.020
成人	2.33	63.2	TEHP	0.100	0.0032
(>18 岁)			TDBPP	—	2.30

2 结果与讨论

2.1 水源水中 OPEs 的总体浓度水平

13 种 OPEs 在长江南京段水源水中的检出情况见表 3, TMP、TEP、TnBP、TBOEP、TCEP、TCPP、

TCrP 和 EHDPP 100% 检出, TPrP 检出率为 7%, TDBPP 无检出。12 种 OPEs 总浓度范围为 $85.21 \sim 1557.96 \text{ ng} \cdot \text{L}^{-1}$, 中值浓度为 $283.80 \text{ ng} \cdot \text{L}^{-1}$, 平均浓度为 $330.88 \text{ ng} \cdot \text{L}^{-1}$, 氯代磷酸酯是最主要化合物, 占总体检出浓度的 57.38%, 远高于烷基磷酸酯(38.50%)和芳香族磷酸酯(4.13%)。

表 3 OPEs 在水源水中的总体检出水平/ $\text{ng} \cdot \text{L}^{-1}$

Table 3 Summary of OPEs concentrations in the sampled source water/ $\text{ng} \cdot \text{L}^{-1}$

化合物	检出率/%	平均值	中值	最大值	最小值
TMP	100	11.15	8.50	55.43	0.70
TEP	100	93.23	55.27	1090.63	4.44
TPrP	7	0.51	ND	12.27	ND
TnBP	100	7.70	7.74	24.94	0.98
TBOEP	100	5.96	3.54	24.18	0.37
TEHP	57	8.82	2.80	144.36	ND
TCEP	100	107.24	44.62	447.08	10.13
TCPP	100	46.74	26.70	318.23	6.02
TDCPP	70	35.87	40.41	264.15	ND
TDBPP	0	ND	ND	ND	ND
TCrP	100	8.70	1.36	170.6	0.22
TPhP	85	0.98	0.96	5.42	0.10
EHDPP	100	3.98	2.33	31.02	0.13
总和		330.88	283.80	1557.96	85.21

长江南京段水源水中平均浓度最高的单体是 TCEP, 浓度范围为 $10.13 \sim 447.08 \text{ ng} \cdot \text{L}^{-1}$, 最高比例为总浓度的 67.93%, TCEP 主要用作塑料制品、聚氨酯泡沫和纺织品的阻燃剂^[4], 有研究发现, TCEP 具有神经毒性、内分泌干扰性、基因突变性和致癌等多种毒性, 已被世界卫生组织(WHO)列为致癌物, 许多发达国家已严格限制 TCEP 在家居用品, 尤其是婴儿用品中的使用^[9,25]。长江流域一带拥有众多企业, 尤其以化工、纺织、服装、电子通信设备、化纤橡胶和塑料等行业为主, 是阻燃剂生产和使用的集聚地, 本研究中 TCEP 检出浓度水平较高, 这可能与工业区仍然在产品中大量添加 TCEP 有关, 有报道指出^[16,26], TCEP 具有较低的 $\lg K_{ow}$ (1.44) 和较高的水溶解度 (7000 mg), 可经空气传输作用于大气环境, 通过干沉降或降雨、降雪等进入河流^[5,27,28], 从而造成水源水中较高的水平。

与其他国家或地区相比, TMP 和 TEP 的高检出率与高浓度在国内外其他河流中很少见^[1,8,29]。西班牙加利西亚的地表水发现 TEP、TCEP、TCPP 和 TnBP 的浓度为 $3 \sim 47 \text{ ng} \cdot \text{L}^{-1}$ ^[30], 德国莱茵河中 OPEs 总浓度为 $10 \sim 200 \text{ ng} \cdot \text{L}^{-1}$ ^[31], 都低于长江南京段水源水中 OPEs, 英国艾尔河中所有样品中均检测到 TCEP、TCPP、TDCPP 和 TPhP, 其中 TCPP 浓度最高, 为 $113 \sim 26050 \text{ ng} \cdot \text{L}^{-1}$, 远高于长江南京段含量水平。与国内部分河流湖泊相比, 本研究总体浓度

水平与太湖($0.82 \sim 1\,066.00 \text{ ng}\cdot\text{L}^{-1}$)^[14], 松花江($5 \sim 3\,700 \text{ ng}\cdot\text{L}^{-1}$)^[15]相当; 汇入渤海的 40 条主要河流中, 11 种 OPEs 范围为 $9.60 \sim 1\,549 \text{ ng}\cdot\text{L}^{-1}$, 平均值为 $300 \text{ ng}\cdot\text{L}^{-1}$, TCP (4.6 ~ 921 $\text{ng}\cdot\text{L}^{-1}$, 平均值: $186 \text{ ng}\cdot\text{L}^{-1}$) 和 TCEP ($1.3 \sim 268 \text{ ng}\cdot\text{L}^{-1}$, 平均值: $80.2 \text{ ng}\cdot\text{L}^{-1}$) 是最丰富的 OPEs, 含氯的 3 种 OPEs (TCP、TCEP、TDCPP) 约占 $\sum$ OPEs 的 91%^[12]; 成都市母亲河——锦江的表层水中 $\sum$ 7OPEs 的浓度范围为 $689.09 \sim 10\,623.94 \text{ ng}\cdot\text{L}^{-1}$ (平均值 $3\,747.58 \text{ ng}\cdot\text{L}^{-1}$), TBOEP 浓度最高, 占 $\sum$ 7OPEs 的质量分数为 36.50% ~ 95.90%^[32]; 北京城市地表水中 14 种 OPEs 浓度为

$3.24 \sim 10\,945 \text{ ng}\cdot\text{L}^{-1}$ (平均值 $954 \text{ ng}\cdot\text{L}^{-1}$)^[8], 比本研究高近 1 个数量级. 从整体上来看, 长江南京段水体中 OPEs 浓度处于较低水平.

2.2 水源水中 OPEs 的时空分布

13 种 OPEs 的时空变异如图 2 所示. 2018 年 8 月 $\sum$ OPEs 的浓度范围为 $220 \sim 1\,557.96 \text{ ng}\cdot\text{L}^{-1}$, 平均值为 $493.78 \text{ ng}\cdot\text{L}^{-1}$, 2018 年 11 月 $\sum$ OPEs 的浓度范围为 $85.21 \sim 354.34 \text{ ng}\cdot\text{L}^{-1}$, 平均值为 $188.31 \text{ ng}\cdot\text{L}^{-1}$, 2019 年 3 月 $\sum$ OPEs 的浓度范围为 $149.11 \sim 446.48 \text{ ng}\cdot\text{L}^{-1}$, 平均值为 $292.74 \text{ ng}\cdot\text{L}^{-1}$, OPEs 浓度随时间呈现明显变化趋势: 8 月 (丰水期) > 3 月 (平水期) > 11 月 (枯水期); 在季节变化方面, 主要污染物浓度为: 夏季 > 春季 > 秋季,




图2 OPEs 在长江南京段水源水中的时空分布与变化

Fig. 2 Spatiotemporal variation of 13 OPEs in source water from the Yangtze River (Nanjing section)

Shi 等^[8]对北京城市地表水(河流和湖泊水)中 14 种 OPEs 的发生,空间分布和季节变化进行了调查,发现 OPEs 浓度因季节而异,夏季 OPEs 水平更高,尤其是 TCPP、TCEP、TMP 和 TEP,这一结果与本研究的结果相似,可见环境水体中 OPEs 浓度水平在不同时期分布特征不同。

11 月(秋季)和 3 月(春季)浓度最高单体是 TEP,分别为 $4.44 \sim 99.47 \text{ ng} \cdot \text{L}^{-1}$ 和 $42.34 \sim 242.39 \text{ ng} \cdot \text{L}^{-1}$ 。8 月(夏季)主要的单体是 TCEP,平均值为 $239.26 \text{ ng} \cdot \text{L}^{-1}$,最高浓度出现在采样点 S4, S4 位于梅山江段,污染可能来源周边较高使用量带来的排放;TEP 为浓度第二高单体,平均浓度为 $106.55 \text{ ng} \cdot \text{L}^{-1}$,其水平与 3 月相当,最高浓度($1090.63 \text{ ng} \cdot \text{L}^{-1}$)出现在 S1,远远高于其他采样点,可能此处存在点源污染。夏季 TCEP、TEP 和 TCPP 这 3 种单体浓度高,TEHP 检出浓度相对低,而 TPrP 仅在 11 月 S14、3 月 S8 和 S11 这 3 个采样点被检出,这表明,可能不同类型的 OPEs 在不同的季节有不同的使用。此外,有研究发现,当温度升高时,OPEs 更易扩散释放到大气中^[7],目前在户外环境中,如空气颗粒^[33]和灰尘^[34,35],已观察到较高水平的 OPEs。南京属亚热带季风气候,四季分明,夏季气温高,雨量

充沛,TCEP、TEP 和 TCPP 容易挥发,更有可能从车辆和建筑中释放出来,扩散到大气中,而空气和灰尘中这些污染物很可能通过湿沉积或降雨释放到河流而富集^[8,9]。相比氯代 OPEs,夏季水体中烷基磷酸酯类可以通过光照或微生物代谢快速分解^[7,26],因此检出浓度相对较低。

2.3 水源水中 OPEs 风险评价

2.3.1 生态风险评估

本研究对长江南京段水源水中 11 种高检出的 OPEs(检出率大于 60%)进行了生态风险评估,从文献[8,13]可知,获得藻类、甲壳类和鱼类的毒理学数据(LC_{50} 或 EC_{50}),评估因子(AF)取 1 000。在水中(表 4),大多数 OPEs 的 RQ 低于 0.1,藻类和鱼类中 RQ 值最高的是 TCrP,分别为 0.588 和 1.55,具有中或高风险,甲壳类中 RQ 值最高的是 EHDPP,值为 1.72,具有高风险。长江南京段水源水中的藻类、甲壳类和鱼类总 RQ 值分别为 $2.70 \times 10^{-3} \sim 0.595$ 、 $3.94 \times 10^{-3} \sim 0.634$ 和 $6.73 \times 10^{-2} \sim 1.56$;这些 OPEs 在水生环境中联合毒性风险较高。Shi 等^[8]在北京地表水中也发现 OPEs 对藻类、甲壳类和鱼类具有较低或中等风险。因此长江南京段水体中的 OPEs 对水生生物和生态环境的风险不可忽视。

表 4 11 种 OPEs 毒理数据及生态风险

Table 4 Toxicity data and ecological risk of eleven OPEs

化合物	敏感物种	$\text{LC}_{50}/\text{mg} \cdot \text{L}^{-1}$	$\text{PNEC}/\text{ng} \cdot \text{L}^{-1}$	RQ
TMP	Fish (<i>Pimephales promelas</i>)	7.0×10^3	7.0×10^6	$9.94 \times 10^{-8} \sim 7.92 \times 10^{-6}$
	Algae (<i>Scenedesmus subspicatus</i>)	9.0×10^2	9.0×10^5	$4.94 \times 10^{-6} \sim 1.21 \times 10^{-3}$
TEP	Crustacean (<i>Daphnia magna</i>)	3.5×10^2	3.5×10^5	$1.27 \times 10^{-5} \sim 3.12 \times 10^{-3}$
	Fish (<i>Leuciscus idus</i>)	2.1×10^3	2.1×10^6	$2.12 \times 10^{-6} \sim 5.19 \times 10^{-4}$
TnBP	Algae (<i>Scenedesmus subspicatus</i>)	4.20	4.2×10^3	$2.33 \times 10^{-4} \sim 5.94 \times 10^{-3}$
	Crustacean (<i>Daphnia magna</i>)	3.65	3.65×10^3	$2.68 \times 10^{-4} \sim 6.83 \times 10^{-3}$
	Fish (<i>Carassius auratus</i>)	5.10	5.1×10^3	$1.92 \times 10^{-4} \sim 4.89 \times 10^{-3}$
TBOEP	Crustacean (<i>Daphnia magna</i>)	75.0	7.5×10^4	$4.90 \times 10^{-6} \sim 3.22 \times 10^{-4}$
	Fish (<i>Pimephales promelas</i>)	13.0	1.3×10^4	$2.83 \times 10^{-5} \sim 1.86 \times 10^{-3}$
TCEP	Algae (<i>Scenedesmus subspicatus</i>)	51.0	5.1×10^4	$1.99 \times 10^{-4} \sim 8.77 \times 10^{-3}$
	Crustacean (<i>Daphnia magna</i>)	3.3×10^2	3.3×10^5	$3.07 \times 10^{-5} \sim 1.35 \times 10^{-3}$
	Fish (<i>Carassius auratus</i>)	90.0	9.0×10^4	$1.13 \times 10^{-4} \sim 4.97 \times 10^{-3}$
TCPP	Algae (<i>Scenedesmus subspicatus</i>)	45.0	4.5×10^4	$1.34 \times 10^{-4} \sim 7.07 \times 10^{-3}$
	Crustacean (<i>Daphnia magna</i>)	91.0	9.1×10^4	$6.61 \times 10^{-5} \sim 3.50 \times 10^{-3}$
	Fish (<i>Poecilia reticulata</i>)	30.0	3.0×10^4	$2.01 \times 10^{-4} \sim 1.06 \times 10^{-2}$
TDCPP	Algae (<i>Scenedesmus subspicatus</i>)	39.0	3.9×10^4	$0 \sim 6.77 \times 10^{-3}$
	Crustacean (<i>Daphnia magna</i>)	4.20	4.2×10^3	$0 \sim 6.29 \times 10^{-2}$
	Fish (<i>Carassius auratus</i>)	5.10	5.1×10^3	$0 \sim 5.18 \times 10^{-2}$
TCrP	Algae (<i>Scenedesmus subspicatus</i>)	0.29	2.9×10^2	$7.55 \times 10^{-4} \sim 0.588$
	Crustacean (<i>Daphnia magna</i>)	0.27	2.7×10^2	$8.11 \times 10^{-4} \sim 0.632$
	Fish (<i>Lepomis macrochirus</i>)	0.11	1.1×10^2	$1.99 \times 10^{-3} \sim 1.55$
TPhP	Algae (<i>Scenedesmus subspicatus</i>)	0.50	5.0×10^2	$0 \sim 1.08 \times 10^{-2}$
	Crustacean (<i>Daphnia magna</i>)	1.00	1.0×10^3	$0 \sim 5.42 \times 10^{-3}$
	Fish (<i>Carassius auratus</i>)	0.70	0.7×10^3	$0 \sim 7.74 \times 10^{-3}$
EHDPP	Crustacean (<i>Daphnia magna</i>)	1.8×10^{-2}	18	$7.35 \times 10^{-3} \sim 1.72$

2.3.2 健康风险评估

本研究评估了长江南京段水源水中 10 种 OPEs 的非致癌风险, 包括 TMP、TEP、TnBP、TBOEP、TEHP、TCEP、TCPP、TDCPP、TPhP 和 TCrP。由于缺乏可用的 SFO 值^[8,23], 仅有 5 种 OPEs (TMP、TnBP、TEHP、TCEP 和 TDCPP) 的致癌风险被讨论。为了更好地评估长江南京段水源水中 OPEs 的最大健康风险, 以每种 OPEs 的最大检出浓度计算 HQ 和 CR。

在最高暴露浓度下, 如图 3 所示, 10 种 OPEs 的总非致癌风险 $\sum HQ$ 为 $4.41 \times 10^{-3} \sim 2.91 \times 10^{-2}$, 均小于 1, 其中 TCEP 非致癌风险最高, HQ 为 1.31×10^{-2} , TPhP 风险最低, HQ 为 2.40×10^{-6} 。且随着年龄的增加, OPEs 的 HQ 逐渐降低, 0~3 岁婴儿 HQ 比成年人高将近一个数量级。

对于致癌风险(图 4), 5 种 OPE 的总致癌风险 $\sum CR$ 值介于 $5.88 \times 10^{-7} \sim 3.89 \times 10^{-6}$ 之间, 其中 TCEP 致癌风险最高, CR 可达 1.84×10^{-6} , TnBP 风险最低, CR 为 6.97×10^{-9} 。根据已有的报

道^[14,20,24], 单个化合物致癌风险的可接受水平为 1.0×10^{-6} , 可知 TCEP 和 TDCPP 对 0~12 个月的儿童有潜在的健康风险, 整体来说, 长江南京段水源水中的 OPEs 对人体的健康风险较低, 与浙江杭州饮用水^[11]、山东潍坊滨海饮用水^[20]、江苏太湖^[13]、骆马湖^[14] 和北京城市地表水^[8] 所报道的健康风险基本一致, 但儿童的健康风险要高于成年人, 需引起高度重视。

3 结论

(1) 长江南京段水源水中共检出了 12 种 OPEs, 总浓度范围为 $85.21 \sim 1557.96 \text{ ng} \cdot \text{L}^{-1}$, TCEP 检出浓度最高, 平均浓度为 $107.24 \text{ ng} \cdot \text{L}^{-1}$, 其次为 TEP 和 TCEP。

(2) OPEs 在长江南京段水源水中呈现明显的时空分布特征, 丰水期 $\sum OPEs$ 浓度明显高于平水期和枯水期; 夏季 $\sum OPEs$ 浓度明显高于春秋季节。

(3) 长江南京段水源水中 TCrP 和 EHDPP 分别对藻类、鱼类和甲壳类有中高生态风险; 高暴露下, TCEP 和 TDCPP 对 0~12 个月的儿童有潜在的健康风险, 应引起相关部门的重视。

参考文献:

- [1] Cristale J, Katsoyiannis A, Sweetman A J, *et al.* Occurrence and risk assessment of organophosphorus and brominated flame retardants in the River Aire (UK) [J]. *Environmental Pollution*, 2013, **179**: 194-200.
- [2] Lee S, Jeong W, Kannan K, *et al.* Occurrence and exposure assessment of organophosphate flame retardants (OPFRs) through the consumption of drinking water in Korea [J]. *Water Research*, 2016, **103**: 182-188.
- [3] 孙佳薇, 丁炜楠, 张占恩, 等. 污水处理厂中有机磷阻燃剂的污染特征 [J]. *环境科学*, 2018, **39**(5): 2230-2238.
Sun J W, Ding W N, Zhang Z E, *et al.* Pollution characteristics of organophosphorus flame retardants in a wastewater treatment plant [J]. *Environmental Science*, 2018, **39**(5): 2230-2238.
- [4] Chen Y Q, Zhang Q, Luo T W, *et al.* Occurrence, distribution and health risk assessment of organophosphate esters in outdoor dust in Nanjing, China: urban vs. rural areas [J]. *Chemosphere*, 2019, **231**: 41-50.
- [5] Li J, Tang J H, Mi W Y, *et al.* Spatial distribution and seasonal variation of organophosphate esters in air above the Bohai and Yellow Seas, China [J]. *Environmental Science & Technology*, 2018, **52**(1): 89-97.
- [6] Wang Y, Sun H W, Zhu H K, *et al.* Occurrence and distribution of organophosphate flame retardants (OPFRs) in soil and outdoor settled dust from a multi-waste recycling area in China [J]. *Science of the Total Environment*, 2018, **625**: 1056-1064.
- [7] 高小中, 许宜平, 王子健. 有机磷酸酯阻燃剂的环境暴露与迁移转化研究进展 [J]. *生态毒理学报*, 2015, **10**(2): 56-68.
Gao X Z, Xu Y P, Wang Z J. Progress in environment exposure, transport and transform of organophosphorus flame retardants [J]. *Asian Journal of Ecotoxicology*, 2015, **10**(2): 56-68.
- [8] Shi Y L, Gao L H, Li W H, *et al.* Occurrence, distribution and seasonal variation of organophosphate flame retardants and plasticizers in urban surface water in Beijing, China [J].




图 3 长江南京段水源水中 10 种 OPEs 的 HQ

Fig. 3 HQ of 10 OPEs in source water from the Yangtze River (Nanjing section)




图 4 长江南京段水源水中 5 种 OPEs 的 CR

Fig. 4 CR of five OPEs in source water from the Yangtze River (Nanjing section)

- Environmental Pollution, 2016, **209**: 1-10.
- [9] 李静, 王俊霞, 许婉婷, 等. 道路灰尘中有机磷阻燃剂污染特征及人体暴露[J]. 环境科学, 2017, **38**(10): 4220-4227.
- Li J, Wang J X, Xu Y T, *et al.* Contamination characteristics and human exposure to organophosphate flame retardants in road dust from Suzhou city[J]. Environmental Science, 2017, **38**(10): 4220-4227.
- [10] Li J, Yu N, Zhang B, *et al.* Occurrence of organophosphate flame retardants in drinking water from China[J]. Water Research, 2014, **54**: 53-61.
- [11] Ding J J, Shen X L, Liu W P, *et al.* Occurrence and risk assessment of organophosphate esters in drinking water from Eastern China[J]. Science of the Total Environment, 2015, **538**: 959-965.
- [12] Wang R M, Tang J H, Xie Z Y, *et al.* Occurrence and spatial distribution of organophosphate ester flame retardants and plasticizers in 40 rivers draining into the Bohai Sea, north China[J]. Environmental Pollution, 2015, **198**: 172-178.
- [13] Liu Y H, Song N H, Guo R X, *et al.* Occurrence and partitioning behavior of organophosphate esters in surface water and sediment of a shallow Chinese freshwater lake (Taihu Lake): implication for eco-toxicity risk[J]. Chemosphere, 2018, **202**: 255-263.
- [14] Xing L Q, Zhang Q, Sun X, *et al.* Occurrence, distribution and risk assessment of organophosphate esters in surface water and sediment from a shallow freshwater Lake, China[J]. Science of the Total Environment, 2018, **636**: 632-640.
- [15] Wang X W, Liu J F, Yin Y G. Development of an ultra-high-performance liquid chromatography-tandem mass spectrometry method for high throughput determination of organophosphorus flame retardants in environmental water[J]. Journal of Chromatography A, 2011, **1218**(38): 6705-6711.
- [16] 刘静, 何丽雄, 曾祥英, 等. 珠江主干和东江河流表层沉积物中有机磷酸酯阻燃剂/增塑剂分布[J]. 生态毒理学报, 2016, **11**(2): 436-443.
- Liu J, He L X, Zeng X Y, *et al.* Occurrence and distribution of organophosphorus flame retardants/plasticizer in surface sediments from the Pearl River and Dongjiang River[J]. Asian Journal of Ecotoxicology, 2016, **11**(2): 436-443.
- [17] 陈政宏, 徐怀洲, 宋宁慧, 等. 高效液相色谱-串联质谱法同时测定水体和沉积物中12种有机磷酸酯类化合物[J]. 分析化学, 2017, **45**(7): 987-995.
- Chen M H, Xu H Z, Song N H, *et al.* Simultaneous determination of 12 kinds of organophosphates in water and sediment by high performance liquid chromatography-tandem mass spectrometry[J]. Chinese Journal of Analytical Chemistry, 2017, **45**(7): 987-995.
- [18] Yadav I C, Devi N L, Li J, *et al.* Concentration and spatial distribution of organophosphate esters in the soil-sediment profile of Kathmandu Valley, Nepal: implication for risk assessment[J]. Science of the Total Environment, 2018, **613-614**: 502-512.
- [19] Li J F, He J H, Li Y, *et al.* Assessing the threats of organophosphate esters (flame retardants and plasticizers) to drinking water safety based on USEPA oral reference dose (RfD) and oral cancer slope factor (SFO)[J]. Water Research, 2019, **154**: 84-93.
- [20] 董政, 马玉龙, 李珺琪, 等. 潍坊滨海经济技术开发区饮用水中有机磷酸酯的水平及人体暴露风险评估[J]. 环境科学, 2017, **38**(10): 4212-4219.
- Dong Z, Ma Y L, Li J Q, *et al.* Occurrence and human exposure risk assessment of organophosphate esters in drinking water in the Weifang Binhai economic-technological Development Area[J]. Environmental Science, 2017, **38**(10): 4212-4219.
- [21] 环境保护部. 中国人群暴露参数手册(成人卷)[M]. 北京: 中国环境出版社, 2013.
- [22] US EPA (Environmental Protection Agency). Guidance on selecting age groups for monitoring and assessing childhood exposures to environmental contaminants[EB/OL]. <https://www.epa.gov/sites/production/files/2013-09/documents/agegroups.pdf>, 2017-06-26.
- [23] Li J F, Zhang Z Z, Ma L Y, *et al.* Implementation of USEPA RfD and SFO for improved risk assessment of organophosphate esters (organophosphate flame retardants and plasticizers)[J]. Environment International, 2018, **114**: 21-26.
- [24] Li W H, Shi Y L, Gao L H, *et al.* Occurrence, distribution and risk of organophosphate esters in urban road dust in Beijing, China[J]. Environmental Pollution, 2018, **241**: 566-575.
- [25] Mizouchi S, Ichiba M, Takigami H, *et al.* Exposure assessment of organophosphorus and organobromine flame retardants via indoor dust from elementary schools and domestic houses[J]. Chemosphere, 2015, **123**: 17-25.
- [26] 杨婷, 何明靖, 杨志豪, 等. 模拟三峡库区消落带土壤有机磷酸酯淹水释放特征[J]. 环境科学, 2018, **39**(12): 5487-5493.
- Yang T, He M J, Yang Z H, *et al.* Simulation of the migration and release characteristics of organophosphate esters in fluctuation zone soil of the Three Gorges reservoir during flooding[J]. Environmental Science, 2018, **39**(12): 5487-5493.
- [27] Cao Z G, Xu F C, Covaci A, *et al.* Distribution patterns of brominated, chlorinated, and phosphorus flame retardants with particle size in indoor and outdoor dust and implications for human exposure[J]. Environmental Science & Technology, 2014, **48**(15): 8839-8846.
- [28] 印红玲, 李世平, 叶芝祥, 等. 成都市大气PM_{2.5}中有机磷阻燃剂的污染水平及来源[J]. 环境科学, 2015, **36**(10): 3566-3572.
- Yin H L, Li S P, Ye Z X, *et al.* Pollution level and sources of organic phosphorus esters in Airborne PM_{2.5} in Chengdu city[J]. Environmental Science, 2015, **36**(10): 3566-3572.
- [29] Cristale J, García Vázquez A, Barata C, *et al.* Priority and emerging flame retardants in rivers: occurrence in water and sediment, *Daphnia magna* toxicity and risk assessment[J]. Environment International, 2013, **59**: 232-243.
- [30] Rodil R, Quintana J B, Concha-Graña E, *et al.* Emerging pollutants in sewage, surface and drinking water in Galicia (NW Spain)[J]. Chemosphere, 2012, **86**(10): 1040-1049.
- [31] Bollmann U E, Möller A, Xie Z Y, *et al.* Occurrence and fate of organophosphorus flame retardants and plasticizers in coastal and marine surface waters[J]. Water Research, 2012, **46**(2): 531-538.
- [32] 吴迪, 印红玲, 李世平, 等. 成都市锦江表层水和沉积物中有机磷酸酯的污染特征[J]. 环境科学, 2019, **40**(3): 1245-1251.
- Wu D, Yin H L, Li S P, *et al.* Pollution characteristics of OPEs in the surface water and sediment of the Jinjiang river in Chengdu city[J]. Environmental Science, 2019, **40**(3): 1245-1251.
- [33] Ren G F, Chen Z, Feng J L, *et al.* Organophosphate esters in total suspended particulates of an urban city in East China[J]. Chemosphere, 2016, **164**: 75-83.
- [34] Khan M U, Li J, Zhang G, *et al.* New insight into the levels, distribution and health risk diagnosis of indoor and outdoor dust-bound FRs in colder, rural and industrial zones of Pakistan[J]. Environmental Pollution, 2016, **216**: 662-674.
- [35] Fromme H, Lahrz T, Kraft M, *et al.* Organophosphate flame retardants and plasticizers in the air and dust in German daycare centers and human biomonitoring in visiting children (LUPE 3)[J]. Environment International, 2014, **71**: 158-163.

CONTENTS

Spatiotemporal Variations of PM _{2.5} Pollution Evolution in China in Recent 20 Years	SHI Yan, LIU Rui-mei, LUO Yi, <i>et al.</i>	(1)
Characteristics of Aerosol Particulate Concentrations and Their Climate Background in Shanghai During 2007-2016	ZHENG Qing-feng, SHI Jun, TAN Jian-guo, <i>et al.</i>	(14)
Characteristics and Sources of Elements in PM _{2.5} During Summer for Three Typical Cities in Pingdingshan-Suizhou-Wuhan, Central China	LIU Wei-jie, SHI Ming-ming, CHENG Cheng, <i>et al.</i>	(23)
PM _{2.5} Pollution Characteristics During Winter and Summer in the Hohhot-Baotou-Ordos Region, China	LI Yi-ming, LIU Jia-yuan, SHI Guo-liang, <i>et al.</i>	(31)
Analysis of Pollution Characteristics and Sources of PM _{2.5} During Heavy Pollution in Shijiazhuang City Around New Year's Day 2019	ZHOU Jing-bo, DUAN Jing-chun, WANG Jian-guo, <i>et al.</i>	(39)
Modeling Studies of Source Contributions to PM _{2.5} in Chengdu, China	XU Yan-ling, YI Ai-hua, XUE Wen-bo	(50)
Chemical Characteristics of Arsenic in PM _{2.5} in Beijing	SHI Shao-xuan, YANG Yan-rong, QIN Juan-juan, <i>et al.</i>	(57)
Pollution Characteristics and Potential Source Contributions of Gaseous Elemental Mercury (GEM) During Summer and Autumn in Ningbo City	YI Hui, LIN Jia-mei, CAI Qiu-liang, <i>et al.</i>	(65)
Seasonal Characteristics and Source Analysis of Water-Soluble Ions in PM _{2.5} of Anyang City	SUN You-chang, JIANG Nan, WANG Shen-bo, <i>et al.</i>	(75)
Analysis of Stable Carbon Isotope Characteristics of PM _{2.5} in Summer and Winter Season in Xinxiang City	YAN Guang-xuan, SONG Xin, ZHANG Jing-wen, <i>et al.</i>	(82)
Construction and Evaluation on Size Resolved Source Apportionment Methods Based on Particle Size Distribution of Chemical Species	LIANG Yong-li, TIAN Ying-ze, LIU Tong, <i>et al.</i>	(90)
Fine Particulate Emission Characteristics of an Ultra-Low Emission Coal-Fired Power Plant	WANG Run-fang, MA Da-wei, JIANG Shao-yi, <i>et al.</i>	(98)
Driving Factors of the Significant Increase in Surface Ozone in the Beijing-Tianjin-Hebei Region, China, During 2013-2018	YU Yi-jun, MENG Xiao-yan, WANG Zhen, <i>et al.</i>	(106)
Characteristic of Surface Ozone and Meteorological Parameters Analysis in Anyang City	YANG Jian, YIN Sha-sha, YU Shi-jie, <i>et al.</i>	(115)
Emission Inventory of Intermediate Volatility Organic Compounds from Vehicles in the Yangtze River Delta in 2017 and the Impact on the Formation Potential of Secondary Organic Aerosols	WANG Qian, HUANG Ling, WANG Yang-jun, <i>et al.</i>	(125)
Exhaust Smoke of Non-Road Vehicles in Typical Cities of Eastern China	LU Jun, HU Qing-yao, YANG Qiang, <i>et al.</i>	(133)
Emission Status and Standards of Volatile Organic Compounds from Chinese and Foreign Bulk Petroleum Terminals	HU Wei, REN Bi-qi, HUANG Yu-hu, <i>et al.</i>	(139)
Secondary Organic Aerosol Mass Yield and Characteristics from 4-ethylguaiaicol Aqueous ·OH Oxidation: Effects of Initial Concentration	ZHUANG Yu, CHEN Yan-tong, LI Xu-dong, <i>et al.</i>	(146)
Stable Isotope Characteristics and Vapor Source of Precipitation in the South and North Slopes of Wushaoling Mountain	XU Xiu-ting, JIA Wen-xiong, ZHU Guo-feng, <i>et al.</i>	(155)
Characteristics and Influencing Factors of Stable Hydrogen and Oxygen Isotopes in Groundwater in the Permafrost Region of the Source Region of the Yangtze River	WANG Shao-yong, HE Xiao-bo, DING Yong-jian, <i>et al.</i>	(166)
Hydrochemical and Isotopic Characteristics of the Lake Balkhash Catchment, Kazakhstan	SHEN Bei-bei, WU Jing-lu, Jilili Abuduwali, <i>et al.</i>	(173)
Effect of a Comprehensive Improvement Project on Water Quality in Urban Lakes: A Case Study of Water Quality Variation in Lihu Lake Over the Past 30 Years	TIAN Wei, YANG Zhou-sheng, SHAO Ke-qiang, <i>et al.</i>	(183)
Detection Method, Distribution, and Risk Assessment of Pharmaceuticals and Personal Care Products in the Yellow Sea and the East China Sea	CHEN Xian, ZHANG Cai-jie, YANG Gui-peng, <i>et al.</i>	(194)
Occurrence and Risk Assessment of Organophosphate Esters in Source Water of the Nanjing Section of the Yangtze River	LI Dong, ZHANG Sheng-hu, ZHANG Qin, <i>et al.</i>	(205)
Spatial Distribution, Spectral Characteristics, and Sources Analysis of Dissolved Organic Matter from Baiyangdian Lake in Xiong'an New District During the Winter Freezing Period	ZHOU Shi-lei, SUN Yue, ZHANG Yi-ran, <i>et al.</i>	(213)
Spatial Characteristics, Risk Assessment, and Source Analysis of Elements in Surface Sediments from the Baiyangdian Lake	WANG Jing-zhong, LIU Zhao, WEI Hao, <i>et al.</i>	(224)
Distribution Characteristics and Ecological Risk Assessment of Phthalate Esters in Surface Sediments of the Songhua River	WANG Huan, YANG Yong-zhe, WANG Hai-yan, <i>et al.</i>	(232)
Occurrence Relationship Between Microplastics and Heavy Metals Pollutants in the Estuarine Sediments of Poyang Lake and the Yangtze River	LI Wen-hua, JIAN Min-fei, LIU Shu-li, <i>et al.</i>	(242)
Effect of Ozonation on Microorganism in the Biological Activated Carbon and Disinfection By-Products in the Effluent	LIU Bao-ming, WANG Xi-xi, ZHANG Xiao-xia, <i>et al.</i>	(253)
Refined Assessment of Exposure and Health Risks of Heavy Metals in Water for the Children in Xigu District, Lanzhou	CHEN Yue-fang, SUN Shan-wei, DUAN Xiao-li, <i>et al.</i>	(262)
Adsorption of Phosphate on Mg/Fe Layered Double Hydroxides (Mg/Fe-LDH) and Use of Mg/Fe-LDH as an Amendment for Controlling Phosphorus Release from Sediments	WU Jun-lin, LIN Jian-wei, ZHAN Yan-hui, <i>et al.</i>	(273)
Characteristics and Mechanism of Monothioarsenate Adsorption on Sand, Sediment, and Goethite	LIAO Dan-xue, SHAN Hui-mei, PENG San-xi, <i>et al.</i>	(284)
Substructure Characteristics of Combined Acid-Base Modified Sepiolite and Its Adsorption for Cd(II)	XIE Sha, XU Ying-ming, YAN Cui-xia, <i>et al.</i>	(293)
Characteristics of Nitrogen Metabolism by Denitrifying Bacterial Agents in Low Dissolved Oxygen Black Odor Water	ZHAO Zhi-rui, LIU Shuo, LI Duo, <i>et al.</i>	(304)
Metagenomic Analysis of Antibiotic Resistant Bacteria and Resistance Genes in a Pharmaceutical and Chemical Wastewater Treatment Plant	CHEN Hong-ling, ZHANG Xing-tao, WANG Qing, <i>et al.</i>	(313)
Measurements of Bacterial Community and Biodiversity from Activated Sludge for a Wastewater Treatment Containing Starch	ZHAO Ting-ting, QIAO Kai, WANG Lei, <i>et al.</i>	(321)
Process Control and Operation Optimization of PN-SAD Coupling Process Based on SBR-ABR	CHEN Chong-jun, ZHANG Min, JIANG Ying, <i>et al.</i>	(330)
Effects of Intermittent Starvation on the Operation of the SNAD Process	LI Dong, LIU Zhi-cheng, XU Gui-da, <i>et al.</i>	(337)
Nitrogen Removal from Mature Landfill Leachate via Denitrification-Partial Nitrification-ANAMMOX Based on a Zeolite Biological Aerated Filter	CHEN Xiao-zhen, WANG Xiao-jun, Karasuta Chayangkun, <i>et al.</i>	(345)
Rapid Achievement of Nitrifying Micro-granular Sludge and Its Nitrification Function	LIU Wen-ru, SONG Jia-jun, WANG Jian-fang, <i>et al.</i>	(353)
Effect of On/Off Aeration Time Ratio Under High Frequency On/Off Aeration on Performance of Nitrosated Granular Sludge	ZHANG Jie, LAO Hui-mei, LI Dong, <i>et al.</i>	(360)
Occurrence, Distribution, Source, and Health Risk Assessment of Polybrominated Diphenyl Ethers in Surface Soil from the Shen-Fu Region, Northeast China	XIANG Xin-xin, LU Yin-tao, RUAN Qi-yang, <i>et al.</i>	(368)
Vertical Pollution Characteristics and Sources of Polycyclic Aromatic Hydrocarbons in a Heterogeneous Unsaturated Zone Under a Coking Plant	MENG Xiang-shuai, WU Meng-meng, CHEN Hong-han, <i>et al.</i>	(377)
Distribution Characteristics of Antibiotic Resistance Genes and Mobile Genetic Elements in Beijing Vegetable Base Soils	ZHANG Ru-feng, SONG Yuan, GAO Hao-ze, <i>et al.</i>	(385)
Stoichiometric Characteristics of Purple Sloping Cropland Under Long-term Fertilization and Cultivation	DU Ying-ni, LI Tian-yang, HE Tian-hui, <i>et al.</i>	(394)
Ecological Stoichiometry of Carbon, Nitrogen, and Phosphorus in Subtropical Paddy Soils	SONG Jia-ling, SHENG Hao, ZHOU Ping, <i>et al.</i>	(403)
Denitrification and Anaerobic Ammonium Oxidation in Soil Nitrogen Migration Process in a Farmland of Wanshangang Lake	CHEN Zhi-hao, QIN Yun-bin, DING Bang-jing, <i>et al.</i>	(412)
Effects of Straw Biochar on Soil Microbial Metabolism and Bacterial Community Composition in Drip-irrigated Cotton Field	WANG Jing, MA Li-juan, LONG Ze-hua, <i>et al.</i>	(420)
Source Appointment of Heavy Metals in Agricultural Soils of the Jiulong River Basin Based on Positive Matrix Factorization	HUANG Hua-bin, LIN Cheng-qi, HU Gong-ren, <i>et al.</i>	(430)
Spatial Distribution of Soil Arsenic and Arsenic Enrichment in Crops in the Oasis Region of the Southeastern Tarim Basin	CHEN Yun-fei, ZHOU Jin-long, ZENG Yan-yan, <i>et al.</i>	(438)
Bioavailability, Translocation, and Accumulation Characteristic of Heavy Metals in a Soil-Crop System from a Typical Carbonate Rock Area in Guangxi, China	MA Hong-hong, PENG Min, LIU Fei, <i>et al.</i>	(449)
Effects of Different Amendments on Fractions and Uptake by Winter Wheat in Slightly Alkaline Soil Contaminated by Cadmium and Nickel	ZHANG Jing-jing, ZHU Shuang-ge, ZHU Li-nan, <i>et al.</i>	(460)
Effect of Different Passivating Agents on the Stabilization of Heavy Metals in Chicken Manure Compost and Its Maturity Evaluating Indexes	LUAN Run-yu, GAO Shan, XU Ying-ming, <i>et al.</i>	(469)
Response of Soil Respiration and Its Components to Nitrogen and Phosphorus Addition in Farming-Withdrawn Grassland in the Semiarid Loess Hilly-Gully Region	ZHOU Jun-jie, CHEN Zhi-fei, YANG Quan, <i>et al.</i>	(479)
Subsequent Effects of Slag and Biochar Application on Greenhouse Gas Emissions from Paddy Fields in the Fuzhou Plain	LAN Xing-fu, WANG Xiao-tong, ZHOU Ya-xin, <i>et al.</i>	(489)